

THE MEDICINAL POTENTIAL OF *ARNEBIA BENTHAMII* IN WOUND HEALING FROM TRADITIONAL USES TO MODERN THERAPEUTICS.

Pooja Pradhan¹, Kamran Javed Naquvi^{2*}, Neeraj Kumar³

^{1,2}Department of Pharmacy, Institute of Biomedical Education and Research Mangalayatan University, Aligarh-202146, India. Mo. 9161504687

³Dr. R.M.L. Institutes of Pharmacy, Badagaon, Shahjahanpur-242401, India

*Corresponding Author:

E-mail address: kjnaquvi@gmail.com

ABSTRACT

Arnebia benthamii, commonly known as Ratanjot, has been traditionally used in herbal medicine for its potent wound-healing properties. This review explores its medicinal potential, tracing its journey from traditional applications to modern therapeutics. Rich in bioactive compounds such as alkannins, flavonoids, and phenolic acids, *A. benthamii* exhibits antimicrobial, anti-inflammatory, and antioxidant properties that accelerate wound healing. Traditional practices have long utilized its extracts for treating cuts, burns, and ulcers, often in the form of ointments or decoctions. Recent pharmacological studies provide scientific validation for these traditional claims, demonstrating the plant's ability to enhance fibroblast proliferation, collagen synthesis, and angiogenesis-key processes in tissue regeneration. Moreover, its antimicrobial activity against wound-associated pathogens underscores its potential as a natural alternative to synthetic antibiotics. Nanotechnology and drug delivery advances have further explored its integration into hydrogel dressings and biomaterials for enhanced wound care. Despite its promising therapeutic applications, challenges such as standardization, bioavailability, and clinical validation remain. Future research should focus on detailed mechanistic studies, clinical trials, and formulation development to harness its full potential in modern medicine. This review highlights *A. benthamii* as a valuable candidate for innovative wound care solutions, bridging the gap between traditional medicine and contemporary pharmacology.

Keywords: *Arnebia benthamii*, wound healing, traditional medicine, phytochemicals, modern therapeutics.

1. Introduction

Wound healing is a complex and dynamic biological process that restores the integrity of damaged skin and tissues. It involves a series of overlapping phases, including haemostasis, inflammation, proliferation, and remodelling [02]. The healing process is influenced by various factors, such as age, nutrition, infection, and underlying medical conditions like diabetes. Proper wound management is crucial to prevent complications like chronic wounds, infections, and excessive scarring. Conventional wound healing treatments include antibiotics, antiseptics, and synthetic dressings. However, these may have limitations, such as antibiotic resistance, delayed healing, or allergic reactions [03]. Hence, natural products and medicinal plants have gained attention as alternative wound healing agents due to their biocompatibility, antimicrobial properties, and regenerative potential. *Arnebia benthamii*, commonly known as Ratanjot or Alkanet, is a valuable medicinal herb native to the Himalayan region. It has been extensively used in traditional medicine systems, such as Ayurveda, Unani, and Tibetan medicine, for treating various ailments, including skin diseases, wounds, burns, and ulcers[01]. The therapeutic significance of *Arnebia benthamii* stems from its bioactive compounds, including alkannins, shikonins, flavonoids, tannins, and phenolic acids, which contribute to its wound-healing, anti-inflammatory, antimicrobial, and antioxidant properties[04]. The root extract of this plant imparts a deep red colour and is often used in herbal formulations for wound treatment and skin regeneration. Despite the extensive traditional use of *Arnebia benthamii*, scientific validation of its wound-healing properties is still in its early stages [05]. Many medicinal plants contain bioactive compounds that facilitate tissue repair, reduce inflammation, and prevent microbial infections. However, systematic studies, including phytochemical characterization, mechanistic insights, and clinical evaluations, are required to establish *Arnebia benthamii* as a reliable therapeutic agent. Given the rise in antibiotic-resistant pathogens and the limitations of synthetic drugs, exploring natural alternatives with broad-spectrum antimicrobial and regenerative potential is essential[06]. *Arnebia benthamii* has demonstrated promising results in preliminary studies, but further research is needed to understand its mechanisms of action, optimize extraction methods, and develop standardized formulations for wound care applications. This review aims to provide a comprehensive analysis of the wound-healing potential of *Arnebia benthamii* by integrating traditional knowledge and modern scientific findings. The key objectives of this review include: Exploring the traditional uses of *Arnebia benthamii* in wound healing and skin-related disorders, Highlighting the phytochemical constituents responsible for its medicinal properties, Examining the mechanisms of action involved in tissue regeneration and antimicrobial defence, Reviewing in vitro, in vivo, and clinical studies that support its therapeutic potential, Discussing current applications, challenges, and future directions in the development of *Arnebia benthamii*-based wound healing formulations, By addressing these aspects, this review aims to bridge the gap between traditional medicine and modern therapeutics, providing insights into the potential integration of *Arnebia benthamii* in natural wound care solutions[11].

Wound healing is a complex biological process that involves multiple cellular and molecular mechanisms, including inflammation, cell proliferation, extracellular matrix remodelling, and tissue regeneration [07]. Impaired wound healing, often associated with infections, chronic inflammation, or underlying health conditions such as diabetes, remains a significant clinical challenge. In recent years, there has been a growing interest in natural remedies derived from medicinal plants, which have historically played a crucial role in traditional medicine. Among these, *Arnebia benthamii*, commonly known as Ratanjot, has been widely recognized for its remarkable wound-healing properties [09]. *A. benthamii*, a member of the Boraginaceae family, is a perennial herb found in the alpine regions of the Himalayas. Traditionally, it has been used in various indigenous healing systems, particularly in Ayurveda and Unani medicine, for treating wounds, burns, and skin infections. The plant's roots are rich in bioactive compounds, including alkannins, shikonins, flavonoids, and phenolic acids, which contribute to its potent antimicrobial, anti-inflammatory, and antioxidant activities. These properties make it a valuable natural agent for promoting wound healing by reducing microbial contamination, enhancing tissue regeneration, and modulating inflammatory responses. In modern therapeutics, the potential of *A. benthamii* is being explored through scientific investigations that validate its pharmacological efficacy. Studies have demonstrated its ability to accelerate fibroblast proliferation, collagen synthesis, and angiogenesis, all of which are critical for effective wound repair [13]. Moreover, advancements in drug delivery systems, such as nanotechnology-based formulations and hydrogel dressings, are being employed to enhance its bioavailability and therapeutic effectiveness. Despite these promising developments, challenges such as standardization, bioavailability, and clinical validation remain key areas of focus. Further research, including in-depth mechanistic studies and clinical trials, is essential to fully harness the medicinal potential of *A. benthamii* in wound care [16]. This review aims to provide a comprehensive overview of its traditional uses, phytochemical composition, and modern therapeutic applications, highlighting its significance as a natural alternative for wound management [17].

2. Traditional Uses of *Arnebia benthamii* in Wound Healing

Arnebia benthamii has been historically documented in traditional medical systems, including Ayurveda, Unani, and Tibetan medicine, for its significant role in skin regeneration and wound healing. In Ayurvedic texts, Ratanjot is recognized for its "Raktashodhak" (blood-purifying) and "Vranaropana" (wound-healing) properties. It has been used in the form of infused oils, herbal pastes, and decoctions to treat burns, ulcers, and surgical wounds [19].

In Unani medicine, *Arnebia benthamii* is classified as having "Mufarre" (anti-inflammatory) and "Mudammil" (wound-healing) properties. It has been traditionally used to prepare Roghan Ratanjot; infused oil applied to wounds and burns for rapid healing and infection prevention [20].

In Tibetan medicine, it has been used as an essential component in herbal formulations to treat skin infections, sores, and chronic wounds. The deep red pigment extracted from the root is known for its cooling and healing effects when applied to damaged tissues[21].

Ethnobotanical Studies: Local and Indigenous Knowledge

Ethnobotanical research has documented the widespread use of *Arnebia benthamii* in Himalayan regions, including India, Nepal, and Tibet, where local communities rely on it for treating skin injuries. Traditional healers prepare pastes and decoctions from its roots and leaves, which are directly applied to cuts, bruises, boils, and ulcers. These communities believe that *Arnebia benthamii* promotes rapid wound contraction, reduces pain, and prevents infections [22].

Traditional Formulations

1. Infused Oils (Roghan Ratanjot)

One of the most popular traditional remedies using *Arnebia benthamii* is Roghan Ratanjot, an oil infusion prepared by soaking the roots in warm oil [25](typically mustard, coconut, or sesame oil). This deep red oil is applied to burns, wounds, and skin infections to:

- Promote faster wound healing
- Reduce inflammation and pain
- Protect against bacterial infections

2. Herbal Decoctions and Pastes

Traditional healers prepare decoctions and herbal pastes by boiling *Arnebia benthamii* roots in water or herbal extracts. The resulting liquid is used for cleansing wounds, while the paste is applied directly to affected areas to enhance collagen synthesis and tissue regeneration.

3. Combination with Other Herbs

Arnebia benthamii is often used in combination with other medicinal herbs to enhance its healing properties:

Turmeric (*Curcuma longa*): Enhances anti-inflammatory and antimicrobial effects

Neem (*Azadirachta indica*): Strengthens antiseptic and antibacterial action.

Aloe Vera: Provides moisturization and soothes burns.

These traditional formulations have been passed down through generations, with strong anecdotal evidence supporting their effectiveness. However, scientific validation through controlled studies is necessary to establish standardized formulations for widespread therapeutic applications.

3. Phytochemical Composition and Bioactive Constituents

Major Chemical Constituents and Their Medicinal Properties are-

Alkannins and Shikonins – Possess strong wound-healing and anti-inflammatory properties, stimulating tissue regeneration and reducing infection risks[23].

Flavonoids – Act as antioxidants that scavenge free radicals and support collagen synthesis, essential for skin repair [27].

Phenolic Compounds – Exhibit antimicrobial and free radical scavenging activities, protecting wounds from oxidative damage and microbial invasion [28].

Tannins and Saponins – Have astringent and antiseptic effects that help in wound contraction and microbial defense [27].

Extraction Techniques for Bioactive Compounds

Various extraction methods are employed to isolate bioactive compounds from *Arnebia benthamii*, including:

- Solvent extraction using ethanol, methanol, or acetone for alkannins and flavonoids.
- Hydrodistillation for isolating essential oils and phenolic compounds.
- Supercritical fluid extraction (SFE) to obtain high-purity extracts with minimal degradation [40].

Comparative Phytochemical Studies of Different Plant Parts

Roots – Richest source of alkannins and shikonins, widely used in wound-healing formulations [29].

S. No.	Chemical Composition and Bioactive Compounds	Uses
1.	Shikonin (acetyl shikonin, isobutyryl shikonin, and β , β -dimethyl acryl shikonin)	Anti-inflammatory, antimicrobial, and antioxidant activities
2.	Alkannins (naphthoquinone compounds)	Wound healing, antimicrobial, and anti-inflammatory effects.
3.	Phenolic Compounds	Antioxidant capacity and overall

		therapeutic potential.
4.	Terpenoids:	Anti-inflammatory and antimicrobial effects.
5.	Essential Oils (Volatile oils)	Antimicrobial and anti-inflammatory
6.	Other Constituents (Fatty Acids, Sterols etc.)	Multifaceted therapeutic profile

Leaves – Contain flavonoids and phenolic acids, exhibiting strong antioxidant properties.

Flowers – Have a moderate concentration of tannins and saponins, contributing to antiseptic action.

These phytochemicals play a pivotal role in the therapeutic effectiveness of *Arnebia benthamii*, supporting its use in traditional and modern wound-healing applications. Future studies should focus on optimizing extraction techniques and developing standardized formulations to enhance its medicinal potential[35].

4. Mechanisms of action in wound healing

Wound healing is a multifaceted biological process essential for restoring skin integrity after injury. It involves a series of overlapping phases: hemostasis, inflammation, proliferation, and remodeling. *Arnebia benthamii*, a medicinal herb widely used in traditional medicine, has garnered attention for its potent wound healing properties [32]. This review focuses on its mechanisms of action, particularly in the modulation of inflammation, promotion of re-epithelialization, enhancement of collagen deposition, stimulation of angiogenesis, and the involvement of molecular pathways and signaling mechanisms[29]. Several studies between 2013 and 2023 have focused on understanding the mechanisms by which *Arnebia benthamii* exerts its wound healing effects. The plant's extract has been demonstrated to influence all four stages of wound healing: hemostasis, inflammation, proliferation, and remodeling. According to Dar et al. (2021), the anti-inflammatory properties of *A. benthamii* help reduce excessive inflammation, which can otherwise delay healing. By modulating pro-inflammatory cytokines like IL-6 and TNF- α , *A. benthamii* helps in creating a conducive environment for wound closure. In addition to its anti-inflammatory effects, *Arnebia benthamii* also promotes cell proliferation and migration, as observed in studies by Mir et al. (2016). The plant extract enhances fibroblast activity and encourages collagen deposition, both of which are crucial for tissue regeneration and wound repair. This has been corroborated by in vivo studies showing that wounds treated with *A. benthamii* extracts heal faster and with less scar formation than untreated wounds (Shafiq et al., 2017)[36].

4.1 Modulation of Inflammation

Inflammation in the skin is the first biological response to tissue injury, which is characterized by the infiltration of immune cells, release of cytokines, and the removal of debris and pathogens [31]. However, excessive or prolonged inflammation can impede the healing process, leading to chronic wounds or excessive scarring. Inhibition of Pro-inflammatory Cytokines: *Arnebia benthamii* modulates the inflammatory phase by downregulating pro-inflammatory cytokines like TNF- α , IL-1 β , and IL-6[37]. The bioactive compound shikonin plays a crucial role in inhibiting the NF- κ B pathway, which is a key regulator of inflammation, thereby reducing the production of these cytokines. Reduction of Nitric Oxide Production: Shikonin also inhibits inducible nitric oxide synthase (iNOS), reducing excessive nitric oxide (NO) levels that can exacerbate inflammation and tissue damage. Promotion of Anti-inflammatory Cytokines: The herb promotes the production of anti-inflammatory cytokines, such as IL-10, aiding in the resolution of inflammation and facilitating the transition to the proliferative phase of wound healing [42].

4.2.2 Promotion of Re-epithelialization

Re-epithelialization is a critical phase in wound healing where new epithelial cells migrate across the wound bed to restore the skin barrier. Keratinocyte Proliferation and Migration: *Arnebia benthamii* promotes the proliferation and migration of keratinocytes, the primary cell type involved in re-epithelialization. Shikonin, through its anti-inflammatory effects, creates a favorable environment for keratinocytes to proliferate and migrate, which is essential for covering the wound bed with new epithelial tissue [47]. Acceleration of Re-epithelialization: Studies have shown that *Arnebia benthamii* accelerates the re-epithelialization process, leading to faster wound closure. This is particularly beneficial in reducing the risk of infection and in minimizing scar formation [48].

4.2.3. Enhancement of Collagen Deposition

Collagen is the main structural protein in the extracellular matrix (ECM) and is crucial for wound strength and integrity. The deposition and remodeling of collagen are key processes during the proliferative and remodeling phases of wound healing. Increased Fibroblast Activity: *Arnebia benthamii* enhances the activity of fibroblasts, the cells responsible for producing collagen [52]. Shikonin stimulates fibroblast proliferation and increases the synthesis of collagen, particularly Type I collagen, which is essential for wound tensile strength. Balanced Collagen Deposition: The herb ensures that collagen deposition is balanced, preventing excessive collagen buildup that can lead to hypertrophic scars or keloids. This balanced deposition is critical for optimal wound healing and functional recovery of the skin[50].

4.2.4. Stimulation of Angiogenesis

Angiogenesis, the formation of new blood vessels, is vital for supplying nutrients and oxygen to the wound site, facilitating tissue repair and regeneration. VEGF Expression: *Arnebia benthamii* stimulates angiogenesis by upregulating Vascular Endothelial Growth Factor (VEGF) expression. VEGF is a key signaling protein that promotes the growth of new blood vessels, ensuring an adequate blood supply to the healing tissue. Enhanced Microcirculation: The herb's ability to enhance microcirculation at the wound site not only speeds up the healing process but also improves the delivery of essential nutrients and oxygen, which are necessary for tissue repair and regeneration [54].

4.2.5. Molecular Pathways and Signaling Mechanisms

The wound-healing effects of *Arnebia benthamii* are mediated through various molecular pathways and signaling mechanisms that orchestrate the complex process of tissue repair [55]. NF- κ B Pathway Inhibition: As mentioned, shikonin inhibits the NF- κ B pathway, which is involved in the inflammatory response. This inhibition reduces the expression of pro-inflammatory cytokines and promotes a timely resolution of inflammation [56]. Activation of TGF- β Pathway: Transforming Growth Factor-beta (TGF- β) is a critical regulator of wound healing, particularly in fibroblast activation and collagen synthesis. *Arnebia benthamii* has been shown to enhance the TGF- β signaling pathway, leading to increased collagen deposition and ECM remodeling [57]. PI3K/Akt Pathway: The PI3K/Akt pathway is another crucial signaling mechanism in wound healing, involved in cell survival, proliferation, and migration. *Arnebia benthamii* activates this pathway, promoting keratinocyte proliferation and migration, as well as fibroblast activity, which are essential for re-epithelialization and collagen deposition. MAPK Pathway: Mitogen-Activated Protein Kinases (MAPKs) play a role in cell proliferation, differentiation, and response to stress. *Arnebia benthamii* modulates the MAPK pathway, enhancing the proliferative and migratory capacity of keratinocytes and fibroblasts, contributing to efficient wound healing[59].

5. CLINICAL STUDIES

Clinical studies on *Arnebia benthamii* are beginning to explore its potential as a therapeutic agent for wound healing. This is based on the promising results of pre-clinical research. This review provides an overview of clinical trials involving *Arnebia benthamii*, its efficacy on different types of ferida, comparisons with treatment with parent ferida, and its safety and tolerability in humans. This is because the majority of research conducted in the past decade has been preclinical. This covers both in vitro and in vivo models. Therefore, there is increasing interest in the clinical use of *Arnebia benthamii* in wound healing. As observed by Khan and colleagues (2022), clinical trials involving topical application of *A. benthamii* in humans have shown promising results. This is especially true in the treatment of chronic burns and furunculosis. Additionally, further clinical trials are needed to validate the therapeutic potential of *A. benthamii* and explore the potential of *A. benthamii* for commercial use in Wound healing products Future research should focus on optimizing the extraction method. Determination of dosage form and monitoring for possible side effects. To guarantee the safe and effective use of this plant in clinical practice. *Arnebia benthamii*, a medicinal herb native to the Himalayan region, has garnered significant attention due to its therapeutic potential, particularly in wound healing[56]. Its roots contain bioactive compounds, such as naphthoquinones (e.g., alkannin and shikonin), which have shown anti-inflammatory, antimicrobial, and antioxidant properties. These properties make it an effective agent in promoting tissue regeneration, reducing inflammation, and preventing infections in wound healing processes. In clinical applications, *Arnebia benthamii* is commonly used in traditional medicine systems like Ayurveda and Unani to treat various skin disorders, wounds, burns, and ulcers. Its extracts are frequently incorporated into topical formulations such as creams, ointments, and oils for external application [60]. Recent studies have also explored its potential in promoting the healing of chronic wounds, diabetic ulcers, and even post-surgical recovery. The therapeutic potential of *Arnebia benthamii* extends beyond wound healing. It also exhibits anti-inflammatory effects, which can help in managing conditions like arthritis, while its antimicrobial properties make it suitable for use in infections and other skin ailments. Moreover, its antioxidant effects contribute to protecting skin from oxidative stress and damage, supporting overall skin health [59].

Clinical Applications and Formulations

Current Pharmaceutical and Cosmetic Applications of *Arnebia benthamii*

Arnebia benthamii has been increasingly incorporated into pharmaceutical and cosmetic products due to its potent wound-healing, anti-inflammatory, and skin-rejuvenating properties[67]. Some of its notable applications include:

- **Topical creams and ointments** for treating burns, ulcers, and cuts.
- **Herbal cosmetics**, including anti-ageing and skin-brightening formulations.
- **Antimicrobial lotions** targeting bacterial and fungal infections.
- **Ayurvedic and Unani medicinal preparations** aimed at promoting tissue regeneration.

Development of Ointments, Gels, and Wound Dressings

Researchers have focused on formulating **novel drug delivery systems** containing *Arnebia benthamii* to enhance its efficacy in wound healing. Some key developments include:

- **Ointments** enriched with *Arnebia benthamii* extract, often combined with **beeswax, turmeric, or neem** to enhance wound recovery.

- **Hydrogel-based formulations**, which provide sustained release of bioactive compounds and maintain moisture for optimal healing.
- **Herbal wound dressings**, infused with *Arnebia benthamii* extracts, offering antimicrobial protection and accelerating tissue repair.

Marketed Herbal Formulations Containing *Arnebia benthamii*

Several herbal and Ayurvedic pharmaceutical companies have commercialized *Arnebia benthamii*-based products, such as:

- **Roghan Ratanjot Oil** – Traditionally used for burns and skin ailments.
- **Herbal wound healing gels** containing *Arnebia benthamii* alongside neem and turmeric.
- **Ayurvedic skin creams and balms**, marketed for their anti-inflammatory and regenerative properties.

Challenges in Standardization, Bioavailability, and Commercialization

Despite its significant medicinal potential, *Arnebia benthamii* faces several challenges in commercial development:

- **Standardization Issues:** Variability in bioactive compound concentrations across different plant sources and extraction methods [66].
- **Low Bioavailability:** Poor water solubility and rapid degradation of key compounds like alkannins and shikonins [64].
- **Regulatory Hurdles:** Need for extensive clinical trials to ensure safety and efficacy [65].
- **Sustainability Concerns:** Overharvesting from the wild leading to the risk of depletion [66].

Addressing these challenges requires **advanced extraction techniques, novel delivery systems (such as nanoparticles), and sustainable cultivation practices** to ensure consistent quality and efficacy of *Arnebia benthamii*-based wound care products [62].

5.1. Overview of two clinical trials involving *Arnebia benthamii*.

Clinical trials involving *Arnebia benthamii* have focused primarily on its application in wound healing. Studies range from small observational trials to more structured randomized clinical trials (RCTs) [68]. These studies examine the effects of wounds on a variety of wound types, including acute, chronic, and surgical wounds. Study Designs: The clinical trials have generally employed topical formulations of *Arnebia benthamii*, such as creams, ointments, or extracts, applied directly to the wound site [69]. The duration of treatment has varied, with most studies monitoring wound healing progress over several weeks. Patient Populations: Trials have included diverse patient populations, including those with diabetic ulcers, pressure sores, burn wounds, and post-surgical wounds, allowing for a broad assessment of the herb's efficacy across different wound types [70].

5.2. Efficacy in Different Types of Wounds

The clinical efficacy of *Arnebia benthamii* has been evaluated across a variety of wound types, with promising results in both acute and chronic wounds. Acute Wounds: In studies involving acute wounds, such as surgical incisions and traumatic injuries, *Arnebia benthamii* has demonstrated a significant reduction in healing time. Patients treated with *Arnebia benthamii*-based formulations experienced faster wound closure, reduced wound size, and minimal scarring compared to those treated with standard care [70]. Chronic Wounds: For chronic wounds, such as diabetic ulcers and pressure sores, *Arnebia benthamii* has shown potential in promoting healing where conventional treatments have often failed [54]. Clinical trials have reported improved wound granulation, enhanced re-epithelialization, and a higher rate of complete wound healing in patients treated with the herb. Surgical Wounds: In post-operative settings, *Arnebia benthamii* has been evaluated for its ability to reduce healing time and prevent complications such as infection and excessive scarring. The results indicate that the herb may be a valuable adjunct to standard surgical wound care, offering faster recovery and improved cosmetic outcomes [49].

5.3. Comparison with Standard Wound Treatments

Arnebia benthamii has been compared to standard wound treatments, such as silver sulfadiazine for burns, saline dressings for ulcers, and other topical antimicrobials. Healing Efficacy: Clinical studies have shown that *Arnebia benthamii* often performs as well as, or better than, standard treatments in terms of wound closure rates, reduction of wound size, and prevention of infection [29]. In some cases, the herb has accelerated healing more effectively than conventional options. Scar Reduction: One of the notable benefits of *Arnebia benthamii* over standard treatments is its ability to minimize scarring. Patients treated with *Arnebia benthamii* have reported better cosmetic outcomes, with less hypertrophic scarring and improved skin texture [49].

5.4. Safety and Tolerability in Human Subjects

Safety and tolerability are crucial aspects of any therapeutic agent. Clinical studies on *Arnebia benthamii* have generally reported favorable safety profiles [23]. Adverse Reactions [44]: The herb has been well-tolerated in most patients, with minimal adverse reactions. Mild skin irritation has been reported in a few cases, but these were typically self-limiting and did not require discontinuation of treatment. Long-term Safety: While short-term use appears safe, long-term safety data is still limited. However, given its traditional use and the absence of serious side effects in clinical trials, *Arnebia benthamii* is considered a low-risk option for wound management. The therapeutic potential of *Arnebia benthamii* in

wound healing and other dermatological applications has led to the development of various formulations. This review covers the types of formulations available, dosage recommendations from studies, and application methods and frequency. Gels: Gels offer a cooling effect and are easily absorbed, making them ideal for burns and inflammatory skin conditions. The gel formulation of *Arnebia benthamii* is favored for its ease of application and patient comfort, especially in cases of painful wounds. Clinical and preclinical studies have provided dosage guidelines to maximize the efficacy of *Arnebia benthamii* formulations in wound healing [21]. The concentration of *Arnebia benthamii* extract in topical formulations typically ranges from 1% to 10%, depending on the severity of the wound and the specific formulation used. Higher concentrations are generally reserved for more severe or chronic wounds. The amount applied per use varies, but studies suggest a thin, even layer of cream, ointment, or gel covering the entire wound area is sufficient. For larger wounds, the amount may need to be adjusted to ensure complete coverage[32].

6. Results and Discussion:

The qualitative phytochemical analysis of extracts from polyherbal drugs has provided valuable insights into the presence and relative concentration of various bioactive compounds, extracted using different solvents: acetone, methanol, ethanol, and chloroform [22]. The solvent extraction efficiency, based on percentage yield, shows ethanol to be the most effective, yield followed by acetone, chloroform, and methanol. This efficiency plays a significant role in optimizing the extraction of potentially therapeutic compounds from plant materials. The results of the present study provide evidence that the antioxidant properties of *Arnebia benthamii* extracts showed mainly the ethyl acetate and ethanol extracts to be the potent source of antioxidants which positively correlates with their total phenolic content [64]. Furthermore, the ethyl acetate and methanol extract also showed potent cytotoxic activity on six human cancer cell lines. Therefore, *Arnebia benthamii* extracts especially ethyl acetate, methanol, and ethanol deserves further investigation in active compounds responsible for the antioxidant and anticancer properties as it might be used in the field of pharmaceutical products and functional foods for the preservation and treatment of cancers[24].

Need for Clinical Trials: Large-Scale Human Studies

Despite promising preclinical and small-scale studies, large-scale human clinical trials are necessary to validate the efficacy and safety of *Arnebia benthamii* in wound healing[23]. Clinical studies should focus on:

- Determining optimal dosage and application frequency.
- Assessing long-term safety and potential side effects.
- Comparing its effectiveness with conventional wound-healing agents.
- Investigating patient-reported outcomes, including pain reduction and cosmetic improvement of scars.

Nanoformulations and Drug Delivery Innovations

To overcome low bioavailability and enhance therapeutic efficiency, researchers are exploring nanoformulations of *Arnebia benthamii*[33]. Nanotechnology-based approaches include:

- **Liposomal carriers**, improving skin penetration and controlled drug release.
- **Nanofiber wound dressings**, offering sustained antimicrobial and anti-inflammatory effects.
- **Nanoemulsions**, increasing the solubility and stability of bioactive compounds like alkannins and shikonins.

These advancements could lead to the development of next-generation wound care products with superior healing properties.

Synergistic Effects with Other Natural Compounds

Combining *Arnebia benthamii* with other herbal and bioactive compounds can potentially enhance wound-healing efficacy. Some promising synergistic combinations include:

- ***Arnebia benthamii* + Turmeric (Curcumin)**: Enhances anti-inflammatory and antimicrobial effects.
- ***Arnebia benthamii* + Aloe Vera**: Promotes hydration, collagen synthesis, and faster tissue regeneration.
- ***Arnebia benthamii* + Honey**: Improves antibacterial properties and accelerates wound closure.

Studying these combinations through rigorous comparative research and mechanistic analysis can help optimize multi-herb formulations for wound management.

Regulatory Challenges and Quality Control Standardization

Despite its therapeutic promise, *Arnebia benthamii* faces regulatory hurdles and quality control challenges that must be addressed for its successful commercialization [11]:

- **Lack of Standardized Extraction Methods**: Variations in bioactive compound concentrations due to differences in geographical origin, processing techniques, and plant maturity [21].
- **Regulatory Approvals**: Need for compliance with global pharmacopoeial standards (e.g., WHO, FDA, EMA) for herbal medicines [52].
- **Batch-to-Batch Consistency**: Establishing protocols for quality control, including HPLC fingerprinting and bioassay-guided standardization [62].
- **Sustainability Concerns**: Overharvesting from wild sources necessitates conservation efforts and large-scale cultivation [42].

Addressing these challenges through advanced analytical methods, regulatory harmonization, and sustainable harvesting practices will pave the way for the global acceptance of *Arnebia benthamii*-based wound healing formulations [32].

7. CONCLUSION:

This study concluded that the ethanolic extract of *Arnebia benthamii* or Ratanjot possesses significant wound healing potential which confers the claim of wound healer plant in Himalayan region. The research data also suggest that *Arnebia benthamii* can be a better choice for polyherbal wound healer preparation. *Arnebia benthamii* is an herb with a long history of use. It shows significant potential in wound healing through its diverse pharmacological properties [10]. Research has highlighted its anti-inflammatory, antimicrobial, and antioxidant effects. This contributes to its effectiveness in promoting wound closure, reducing inflammation, and improving tissue regeneration. Pre-clinical studies show that *Arnebia benthamii* accelerates wound healing by modulating the inflammatory response. Promote the creation of new cells Increase the accumulation of collagen and stimulate the creation of new blood vessels Clinical studies are beginning to examine these effects in human volunteers. It has shown efficacy in the treatment of acute, chronic and surgical ferrida, with a favourable safety profile [31]. In conclusion, *Arnebia benthamii* holds great promise as a natural and available ferrida treatment agent. efficiency with continuous research and development It therefore has the potential to become a fundamental component in modern fever treatment. It offers patients a safe and effective alternative to traditional treatments. Continued exploration of *Arnebia benthamii*'s properties and uses will be fundamental to unlocking its full healing potential. and its application in general clinical practice. In summary, research over the past decade has strengthened *Arnebia benthamii* as a potential wound-healing agent. Because it has anti-inflammatory, antimicrobial, and antioxidant properties. and the creation of new tissue the plant's intense phytochemical properties especially the high chiconine content [43]. It is considered a supporting factor for its treatment effect. As interest in natural and alternative treatments grows, *A. benthamii* holds great promise for integration into modern wound healing approaches. *Arnebia benthamii* (commonly known as *Indian arnebia* or *Torma*) has garnered increasing attention in modern wound healing research due to its potential therapeutic properties. Bridging traditional knowledge with scientific validation, this plant has a long history of use in indigenous medicine, particularly in wound care and skin conditions. Its active compounds, notably alkannin and shikonin (naphthoquinones), have shown promising effects in promoting wound healing, and anti-inflammatory, antioxidant, and antimicrobial activities.

Traditional Knowledge

In traditional systems of medicine like Ayurveda and Traditional Chinese Medicine (TCM), *Arnebia benthamii* has been used for centuries for treating various skin ailments such as wounds, burns, ulcers, and infections. The plant's root extracts have been topically applied to promote healing and reduce scarring. These practices have been passed down through generations, providing a wealth of anecdotal evidence for its efficacy [42].

Scientific Validation

Recent research has substantiated the wound-healing properties of *Arnebia benthamii* through various scientific studies[55]. The key active compounds—**alkannin** and **shikonin**—are found to possess:

- **Anti-inflammatory properties:** Reducing inflammation at the wound site, which is crucial for promoting tissue repair[43].
- **Antioxidant activities:** Protecting cells from oxidative stress and damage, which could otherwise impair healing.
- **Antimicrobial effects:** Helping to prevent infections in open wounds by inhibiting the growth of harmful microorganisms.
- **Promotion of collagen synthesis:** Essential for tissue regeneration and the formation of new skin at the wound site.
- **Increased angiogenesis:** Supporting the formation of new blood vessels, which is vital for proper nutrient and oxygen supply to the healing tissue.

These effects have been observed in both in vitro (lab) studies and in vivo (animal) models, suggesting that *Arnebia benthamii* may be an effective alternative or adjunct in modern wound management [56].

Future Implications for Phytomedicine and Dermatology

1. **Integration into Modern Therapies:** With increasing interest in natural products, *Arnebia benthamii* could be incorporated into commercial wound care formulations, such as creams, gels, and ointments. Its combination of anti-inflammatory, antioxidant, and antimicrobial properties makes it an ideal candidate for topical treatments [46].
2. **Nanotechnology:** Advances in nanotechnology may improve the bioavailability and targeted delivery of *Arnebia benthamii*'s active compounds. Nanoparticles or nano-emulsions can enhance the plant's effectiveness in wound healing by improving its absorption and sustained release at the site of injury [32].
3. **Personalized Phytomedicine:** The growing trend toward personalized medicine might see the use of *Arnebia benthamii*-based products tailored to individual wound healing needs, taking into account the severity of the wound, the patient's skin type, and any underlying conditions [23].
4. **Research in Dermatology:** Beyond wound healing, further research could explore the plant's role in treating chronic dermatological conditions, such as eczema, psoriasis, and acne, due to its anti-inflammatory and skin-regenerative effects.

In conclusion, the convergence of traditional knowledge with modern scientific research opens the door to utilizing *Arnebia benthamii* as a natural agent in wound healing. Its future in phytomedicine and dermatology looks promising, potentially leading to more effective, safer, and holistic wound care treatments. Further clinical studies are needed to confirm its efficacy and safety, but the potential is undeniable [42].

REFERENCES:

1. Ghosh S, Kaur S. Phytochemical composition and traditional uses of *Arnebia benthamii*. J Ethnopharmacol. 2018;210:125-34. doi:10.1016/j.jep.2017.08.034.
2. Thakur M, Kumar A. Chemical constituents of *Arnebia benthamii*: A review. Phytochem Rev. 2020;19(3):475-90. doi:10.1007/s11101-019-09640-2.
3. Jabeen F, Khan M. Antimicrobial properties of *Arnebia benthamii* extracts. Asian Pac J Trop Med. 2019;12(6):271-6. doi:10.4103/1995-7645.258970.
4. Kumar V, Verma R. Anti-inflammatory activity of *Arnebia benthamii* in experimental models. J Herb Med. 2021;23:100-6. doi:10.1016/j.hermed.2020.100406.
5. Sharma S, Gupta P. Efficacy of *Arnebia benthamii* ointment in wound healing: A randomized controlled trial. J Wound Care. 2022;31(4):200-7. doi:10.12968/jowc.2022.31.4.200.
6. Mehta R, Joshi R. Ethnomedicinal uses of *Arnebia benthamii* in Indian traditional medicine. J Ethnopharmacol. 2017;205:152-8. doi:10.1016/j.jep.2017.05.037.
7. Das S, Banerjee A. Mechanistic insights into wound healing properties of *Arnebia benthamii*. BMC Complement Med Ther. 2019;19(1):102. doi:10.1186/s12906-019-2542-7.
8. Singh M, Kumar N. Role of phytochemicals in wound healing: Focus on *Arnebia benthamii*. Pharmacogn Rev. 2020;14(28):50-8. doi:10.4103/phrev.phrev_27_19.
9. Prakash P, Yadav S. In vivo assessment of wound healing potential of *Arnebia benthamii* using animal models. J Nat Prod. 2021;84(2):337-45. doi:10.1021/acs.jnatprod.0c01017.
10. Khan A, Singh R. Comparative analysis of wound healing effects of *Arnebia benthamii* and other herbal remedies. Int J Herb Med. 2020;8(1):28-34.
11. Kumar, S., & Roy, P. (2020). An overview of herbal remedies for wound healing: Spotlight on *Arnebia benthamii*. *Journal of Herbal Medicine*, 21, 100-108. <https://doi.org/10.1016/j.hermed.2020.100408>
12. Joshi H, Shah N. Toxicological profile of *Arnebia benthamii*: Safety evaluation in animal models. J Ethnopharmacol. 2019;243:112-9. doi:10.1016/j.jep.2019.112119.
13. Agarwal S, Kumar R. Ethnobotanical significance and pharmacological properties of *Arnebia benthamii*. Indian J Tradit Knowl. 2021;20(2):365-75.
14. Zubair M, Iqbal Z. The role of *Arnebia benthamii* in regenerative medicine: A comprehensive review. J Regen Med. 2020;15(3):205-15.
15. Hussain A, Shah S. Investigating the antioxidant activity of *Arnebia benthamii* extracts and their role in wound healing. Phytochem Anal. 2022;33(1):78-85. doi:10.1002/pca.3043.
16. Rani A, Sethi J. Wound healing properties of *Arnebia benthamii* and its bioactive compounds. J Med Plants Res. 2019;13(10):250-8. doi:10.5897/JMPR2019.6792.
17. Kaur R, Singh P. Mechanisms of action of *Arnebia benthamii* in cutaneous wound healing. Int J Cosmet Sci. 2021;43(5):465-72. doi:10.1111/ics.12683.
18. Chaudhary S, Bharti S. Synergistic effects of *Arnebia benthamii* with other herbs on wound healing. J Herb Pharmacother. 2020;20(2):93-104.
19. Malik A, Kumar D. Histopathological analysis of wound healing with *Arnebia benthamii* extracts. J Pathol. 2019;239(4):459-67.
20. Yadav V, Joshi P. Antimicrobial and wound healing activities of *Arnebia benthamii* extracts: An in vitro study. Pharm Biol. 2022;60(1):712-8. doi:10.1080/13880209.2022.2027528.
21. Singh R, Kumar R. A review of traditional uses of *Arnebia benthamii* in wound healing. Anc Sci Life. 2018;37(1):50-5. doi:10.4103/asl.ASL_256_17.
22. Verma R, Sharma N. Role of *Arnebia benthamii* in chronic wound management: An evidence-based review. J Wound Care. 2021;30(10):560-7. doi:10.12968/jowc.2021.30.10.560.
23. Pandey A, Chaudhary N. Evaluating the efficacy of *Arnebia benthamii* in diabetic wound healing. Diabetes Metab Syndr. 2020;14(6):2019-24. doi:10.1016/j.dsx.2020.09.033.
24. Kumar H, Gupta A. The impact of *Arnebia benthamii* on collagen synthesis in wound healing. J Cosmet Dermatol. 2019;18(5):1392-9. doi:10.1111/jocd.12983.
25. Bhat S, Ahmad M. Clinical implications of *Arnebia benthamii* for enhancing wound healing. Int J Med Sci. 2022;19(8):1651-9.
26. Singh A, Varma R. Phytotherapeutic approaches using *Arnebia benthamii* for wound healing. J Herb Med. 2021;27:100415. doi:10.1016/j.hermed.2021.100415.
27. Shukla S, Sharma P. Anti-inflammatory properties of *Arnebia benthamii* in wound healing applications. Asian Pac J Trop Med. 2019;12(4):203-10.
28. Gupta R, Meena M. Role of herbal medicine in wound care: A study on *Arnebia benthamii*. J Wound Care Manag. 2020;4(2):150-5.
29. Kumar N, Shukla S. Natural products for wound healing: Spotlight on *Arnebia benthamii*. Pharmacogn J. 2021;13(4):843-50. doi:10.5530/pj.2021.13.105.
30. Joshi S, Yadav A. Synergistic action of *Arnebia benthamii* and silver nanoparticles in wound healing. Mater Sci Eng C. 2018;93:325-33. doi:10.1016/j.msec.2018.08.057.

31. Sahu R, Chatterjee A. The healing effects of *Arnebia benthamii* on excisional wounds in diabetic rats. *J Diabetes Res.* 2022;2022:1-9. doi:10.1155/2022/3475912.
32. Raghav S, Kaur N. Comprehensive analysis of *Arnebia benthamii* as a wound healing agent. *J Nat Remedies.* 2020;20(3):22-30.
33. Sharma A, Jain P. Antioxidant and anti-inflammatory effects of *Arnebia benthamii* in wound healing. *J Ethnopharmacol.* 2021;271:113917. doi:10.1016/j.jep.2021.113917.
34. Ali M, Khan F. The role of *Arnebia benthamii* in modern wound management: A systematic review. *J Wound Care.* 2020;29(5):222-8. doi:10.12968/jowc.2020.29.5.222.
35. Mukherjee P, Kumar S. Assessment of wound healing properties of *Arnebia benthamii* in a murine model. *Phytother Res.* 2022;36(1):15-24. doi:10.1002/ptr.7186.
36. Kapoor R, Verma S. Herbal formulations for wound healing: Focus on *Arnebia benthamii*. *Int J Herb Med.* 2019;7(3):95-100.
37. Gupta A, Chaudhary A. Efficacy of *Arnebia benthamii* in the healing of thermal burns. *J Burn Care Res.* 2020;41(3):453-9. doi:10.1093/jbcr/irz231.
38. Bhattacharya S, Dey A. Comparative analysis of wound healing efficacy of *Arnebia benthamii* and synthetic agents. *Int J Pharmacogn Phytochem Res.* 2021;13(2):300-6.
39. Joshi K, Sharma D. Phytochemical screening and evaluation of *Arnebia benthamii* for its wound healing potential. *J Med Chem.* 2022;65(8):3564-72. doi:10.1021/acs.jmedchem.1c01073.
40. Kumar P, Sharma R. The influence of *Arnebia benthamii* on fibroblast activity in wound healing. *J Cell Sci Ther.* 2018;9(4):345-52.
41. Singh N, Tiwari P. The impact of *Arnebia benthamii* on angiogenesis during wound healing. *J Vasc Biol.* 2020;12(1):45-52.
42. Ahuja A, Saini S. Wound healing effects of *Arnebia benthamii* in excisional and incision models. *Asian J Pharm Clin Res.* 2019;12(6):125-30.
43. Kumar V, Mehta A. Healing properties of *Arnebia benthamii*: A study on its effects on granulation tissue formation. *BMC Complement Med Ther.* 2021;21(1):67. doi:10.1186/s12906-021-03245-2.
44. Nair S, Raghavan S. Immunomodulatory effects of *Arnebia benthamii* in wound healing. *Phytomedicine.* 2020;68:153168. doi:10.1016/j.phymed.2020.153168.
45. Sharma R, Gupta S. Exploring the efficacy of *Arnebia benthamii* in diabetic wound healing. *Diabetes Metab J.* 2022;46(2):150-9. doi:10.4093/dmj.2021.0241.
46. Yadav S, Sharma N. Mechanisms of action of *Arnebia benthamii* in wound healing: A review. *Herb Med Open Access.* 2019;5(1):1-7.
47. Verma P, Bhatia R. Role of *Arnebia benthamii* in skin regeneration and wound healing. *J Tissue Eng Regen Med.* 2021;15(10):861-70. doi:10.1002/term.3147.
48. Mehta S, Kapoor R. Comparative efficacy of *Arnebia benthamii* and synthetic dressings in wound healing. *J Wound Care Manag.* 2022;5(3):230-6.
49. Singh M, Bharti S. Antimicrobial and healing properties of *Arnebia benthamii*: Implications in wound care. *J Nat Remedies.* 2020;20(4):99-106.
50. Singh J, Dhillon S. The therapeutic potential of *Arnebia benthamii* in skin disorders and wound healing. *Pharmacogn Rev.* 2019;13(26):55-63. doi:10.4103/phrev.phrev_13_18.
51. Kumar S, Sinha R. The impact of *Arnebia benthamii* on collagen deposition in wound healing. *J Biomed Sci Eng.* 2021;14(1):41-50.
52. Verma A, Joshi R. Evaluation of topical formulations of *Arnebia benthamii* for enhanced wound healing. *Int J Pharm Pharm Sci.* 2020;12(5):115-20.
53. Patil D, Kulkarni A. Traditional uses and modern applications of *Arnebia benthamii* in wound healing. *J Ayurveda Integr Med.* 2021;12(3):430-8. doi:10.1016/j.jaim.2020.10.008.
54. Akhtar MS, et al. Phytochemical and pharmacological investigations of *Arnebia benthamii* for wound healing activity. *J Ethnopharmacol.* 2014;153(1):136-42. doi:10.1016/j.jep.2013.11.023
55. Wani TA, Dhar SA, Bazaz MR, et al. Evaluation of the wound healing potential of ethanolic extract of *Arnebia benthamii* in rats. *Phytomedicine.* 2013;20(4):359-365. doi:10.1016/j.phymed.2012.12.006
56. Sharma R, Singh R, Gupta R, et al. Wound healing potential of *Arnebia benthamii*: An insight into its traditional use. *Planta Med.* 2016;82(12):1037-1045. doi:10.1055/s-0042-109841
57. Khan SA, Lone AH, Malik SA, et al. Bioactive components and wound healing efficacy of *Arnebia benthamii* root extracts. *J Med Plants Res.* 2015;9(7):219-225. doi:10.5897/JMPR2015.5867
58. Firdous SM, Sofi AH, Zargar MA, et al. Exploring the therapeutic potential of *Arnebia benthamii* in wound healing: In vitro and in vivo studies. *Biomed Pharmacother.* 2017;88:952-960. doi:10.1016/j.biopha.2017.01.028
59. Dandagi PM, Angadi UB, Ravikumar BV, et al. Herbal formulation of *Arnebia benthamii* for wound healing: A pharmacological review. *Indian J Pharm Sci.* 2012;74(2):122-129. doi:10.4103/0250-474X.103850
60. Mahapatra A, Rout SD, Acharya S, et al. Ethnopharmacological evaluation of *Arnebia benthamii* for wound healing properties. *J Herb Med.* 2019;18:100281. doi:10.1016/j.hermed.2019.100281
61. Koul B, Sanyal I, Mehta P, et al. Recent pharmacological and phytochemical aspects of *Arnebia benthamii*: A review. *J Pharm Bioallied Sci.* 2017;9(4):235-242. doi:10.4103/0975-7406.215352

62. Wani K, Nawchoo IA. Traditional uses and phytochemistry of *Arnebia benthamii*: A plant with significant wound healing potential. *J Ayurvedic Herb Med.* 2015;2(4):134-142. doi:10.4103/2348-0365.159882
63. Kaithwas G, Mukherjee A, Majumdar DK. Potential of *Arnebia benthamii* in enhancing wound healing and anti-inflammatory activity. *J Pharm Pharmacol.* 2016;68(9):1189-1200. doi:10.1111/jphp.12574
64. Ahmad S, Hassan MA, Ansari SH. Pharmacological evaluation of *Arnebia benthamii* extracts for their wound healing potential in experimental models. *J Ethnopharmacol.* 2021;275:114061. doi:10.1016/j.jep.2021.114061
65. Bhat ZA, Khan NA, Khuroo MA. Traditional uses and pharmacological potential of *Arnebia benthamii*: A comprehensive review. *Phytomedicine.* 2019;23(7):734-742. doi:10.1016/j.phymed.2019.02.024
66. Rasool B, Habib M, Masoodi MA. Phytochemical constituents and wound healing efficacy of *Arnebia benthamii* extracts. *J Tradit Complement Med.* 2020;10(4):425-432. doi:10.1016/j.jtcme.2019.11.004
67. Mir MA, Koul B, Tiku AB. *Arnebia benthamii*: An underutilized medicinal herb with high potential for wound healing and anti-inflammatory applications. *Pharmacogn Rev.* 2018;12(24):215-223. doi:10.4103/phrev.phrev_25_18
68. Ali SR, Qazi PH, Zargar M. Evaluation of antimicrobial and antioxidant activities of *Arnebia benthamii* in the context of wound management. *Int J Pharm Sci.* 2016;8(6):350-356. doi:10.22377/ijp.v8i6.1028
69. Tabasum H, Shah A, Sofi AH. Role of *Arnebia benthamii* in accelerating wound repair: A systematic review. *J Herb Med.* 2017;7(3):45-52. doi:10.1016/j.hermed.2017.02.005
70. Khan NU, Lone BA, Rather GM. Ethnopharmacological aspects of *Arnebia benthamii* and its relevance in wound healing therapies. *J Ethnobiol Ethnomed.* 2022;18(1):101. doi:10.1186/s13002-022-00494-y
71. Sharma SP, Gupta A. Phytochemical profiling and wound healing efficacy of *Arnebia benthamii* in traditional medicine. *Evid Based Complement Alternat Med.* 2020;2020:687954. doi:10.1155/2020/6879543